Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.



MAR FOOD ADMINISTRATION OFFICE OF MARKETING SERVICES COTTON AND FIBER BRANCH

ESTIMATED GRADE AND STAPLE LENGTH OF AMERICAN-EGYPTIAN (SXP) COTTON GINNED JAN 1 352 BRAR RECEIVED

JANUARY 23, 1945

ALL LENGTHS	LENGTH (INCHES) 1-9/32" AND SHORTER 1-5/16" AND 1-11/32" 1-7/16" AND 1-15/32" 1-9/16" AND 1-17/32" 1-9/16" AND LONGER	ALL GRADES	GRADE AND STAPLE LENGTH 1-1/2 2-1/2 3-1/2 3-1/2 4-1/2 5 AND BELOW
0.001 0.001	25. *0 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -	0.001 0.001	DECEMBER 13 JANUARY 15 JANUARY 15 PCT. PCT. PCT. PCT. PCT. PCT. PCT. PCT.
100.0 100.0	50.25 40.27 50.25 40.27 50.25 40.27	0.001 0.001	ARIZONA-CALIFORNIA SEASON THRU JANUARY 15 JANUARY 15 JANUARY 15 A.C. 36.5 18.6 1.2 28.2 18.6 1.6 1.6 1.8 1.2 1.8 1.2 1.8 1.8 1.8 1.8
1	1 1 1 1 1 1	•	DECEMBER 13 JANUARY 15 1943 1944 PCT. PCT. SSAMPLES
	11111	1	NEW MEXICO II SEASON THRU JANUARY 15 1943 1944 PCT. PCT.
100.0 100.0	49.3 50.5 57.4 57.4	0.001 0.001	TEXAS . 1 DECEMBER 13 JANUARY 15 1943 1944 PCT
0.001 0.001	26.9	0.001 0.001	SEASON THRU JANUARY 15 1943 1944 PCT. PCT. 16.3 20.7 52.0 24.8 14.9 28.4 8.5 15.5 5.7 7.8 1.6 8 2.4 8 2.6 4 8.6 8 2.6 4 8.6 8 2.6 4 8.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.6 8 2.7 8 2.8 8 2.
0.00 100.0	22 *2 *3 *49.4 *49.4 *50.3 *49.4 *49.4 *49.4 *49.4 *49.4 *49.4 *49.4 *49.4 *49.4 *40.6 *40	0.001 0.001	DECEMBER 13 JANUARY 15 1943 1944 PCT. PCT. 8.0 2.4 28.3 11.1 32.7 21.3 6.3 18.2 1.9 20.3 7.4
100.0 100.0	55.2.1. *- 55.2.1. *- 69.3.1.1	0.001 0.001	TAI SEASON THRU JANUARY 15 1943 1944 PCT. PCT. 11.5 13.0 21.2 28.5 20.4 14.0 13.0 15.0 13.0 1.4 7.3 2.1

^{*} LESS THAN 0.05 PERCENT.

1) NO SAMPLES OF AMERICAN-EGYPTIAN (SXP) COTTON HAVE BEEN CLASSED FROM NEW MEXICO THIS SEASON.

WAR FOOD ADMINISTRATION OFFICE OF WARKETING SERVICES COTTON AND FIBER BRANCH BAKERSFIELD, CALIFORNIA

VOLUME XVIII, No. 8

JANUARY 23, 1945

COTTON QUALITY REPORT FOR GINNINGS THROUGH JANUARY 15, 1945

AMERICAN-EGYPTIAN COTTON (SXP)

AMERICAN-EGYPTIAN (SXP) COTTON GINNED THROUGH THE MIDDLE OF JANUARY WAS LOWER IN GRADE BUT NOTICE—ABLY LONGER IN STAPLE THAN GINNINGS TO THE SAME DATE A YEAR AGO, ACCORDING TO THE WAR FOOD ADMINISTRATION. ABOUT 68 PERCENT OF THE CURRENT CROP IS COMPRISED OF GRADES 2-1/2 AND HIGHER AND APPROXIMATELY 32 PERCENT WAS GRADE 3 AND LOWER. LAST SEASON THESE GRADE CATEGORIES WERE REPRESENTED BY PERCENTAGES OF ABOUT 92 AND 8 PERCENT RESPECTIVELY WHICH CONSTITUTES QUITE A LOWERING OF GRADES IN THIS SEASON'S CROP. STAPLES 1-1/2" AND LONGER ACCOUNTED FOR ALMOST 70 PERCENT OF CURRENT GINNINGS THROUGH JANUARY 15 AGAINST ONLY ABOUT 57 PERCENT TO THE SAME DATE LAST YEAR.

ARIZONA—CALIFORNIA - AMERICAN—EGYPTIAN (SXP) GINNINGS IN ARIZONA—CALIFORNIA THROUGH JANUARY 15 WERE LOWER IN GRADE AND SOMEWHAT LONGER IN STAPLE THAN GINNINGS THROUGH THE MIDDLE OF JANUARY

WERE LOWER IN GRADE AND SOMEWHAT LONGER IN STAPLE THAN GINNINGS THROUGH THE MIDDLE OF JANUARY
LAST SEASON. ALMOST 46 PERCENT OF THE CURRENT CROP WAS GRADE 3 AND LOWER AND 58 PERCENT WAS
1-1/2" AND LONGER. LAST YEAR ONLY 9 PERCENT WAS GRADE 3 AND LOWER AND 52 PERCENT WAS 1-1/2" AND
LONGER.

TEXAS - DISTRICT 6 - THE CROPE OF AMERICAN-FOURTLAN (SYP) CLANSED IN DISTRICT 6 OF TEXAS THROUGH

TEXAS - DISTRICT 6 - THE GRADE OF AMERICAN-EGYPTIAN (SXP) GINNED IN DISTRICT 6 OF TEXAS THROUGH
JANUARY 15, 1945 WAS DECIDEDLY LOWER THAN LAST SEASON. GRADES 1 AND 1-1/2 COMPRISED ALMOST 46

PERCENT OF GINNINGS THIS SEASON AGAINST 68 PERCENT A YEAR AGO AND ABOUT 44 PERCENT WAS GRADES 2

AND 2-1/2 COMPARED WITH ONLY 23 PERCENT LAST SEASON. THE PROPORTION OF GRADES 3 AND LOWER WAS

ALMOST 11 PERCENT THIS YEAR AGAINST ABOUT 8 PERCENT LAST YEAR. ALMOST 87 PERCENT OF THE CURRENT

CROP WAS ESTIMATED TO BE 1-1/2" AND LONGER COMPARED WITH ONLY ABOUT 73 PERCENT OF THE GINNINGS

LAST YEAR.

According to the Bureau of the Census, Ginnings through January 15, 1945 WERE: ARIZONA-CALIFORNIA - 4433 BALES; TEXAS - 2996 BALES.